

Octonions

and Octonionic Projective Geometry

Tijana Šukilović

Department of Geometry


Faculty of Mathematics

University of Belgrade, Serbia

Introduction

- William Rowan Hamilton - *quaternions*
- John T. Graves - *octaves*
- Arthur Cayley - *Cayley's numbers*
- Élie Cartan - *geometrical relevance*
- Jordan, von Neumann and Wigner - *the foundations of quantum mechanics*

Inventing the Quaternions



Brougham Bridge - the bridge where Hamilton carved his definition of the quaternions

Octonions: if 4, why not more?

Graves: If you're allowed make up a way of multiplying lists of 4 numbers, why not more?

Hamilton: Just because it's bigger I don't know if it is better. I have a horse with four legs, I don't know if your horse with eight legs will run twice as fast!



Basic Definitions

- A *vector space* is a finite-dimensional module over the field of real numbers.

Basic Definitions

- A **vector space** is a finite-dimensional module over the field of real numbers.
- An **algebra** \mathcal{A} is a vector space that is equipped with a bilinear map $m : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ called ‘multiplication’ and a nonzero element $1 \in \mathcal{A}$ called the ‘unit’ such that $m(1, a) = m(a, 1) = a$.

Basic Definitions

- A **vector space** is a finite-dimensional module over the field of real numbers.
- An **algebra** \mathcal{A} is a vector space that is equipped with a bilinear map $m : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ called ‘multiplication’ and a nonzero element $1 \in \mathcal{A}$ called the ‘unit’ such that $m(1, a) = m(a, 1) = a$.
- An algebra \mathcal{A} is a **division algebra** if given $a, b \in \mathcal{A}$ with $ab = 0$, then either $a = 0$ or $b = 0$.

Basic Definitions

- A **vector space** is a finite-dimensional module over the field of real numbers.
- An **algebra** \mathcal{A} is a vector space that is equipped with a bilinear map $m : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ called ‘multiplication’ and a nonzero element $1 \in \mathcal{A}$ called the ‘unit’ such that $m(1, a) = m(a, 1) = a$.
- An algebra \mathcal{A} is a **division algebra** if given $a, b \in \mathcal{A}$ with $ab = 0$, then either $a = 0$ or $b = 0$.
- A **normed division algebra** is an algebra \mathcal{A} that is also a normed vector space with $\|ab\| = \|a\|\|b\|$.

Levels of Associativity

An algebra is:

- ***power-associative*** if the subalgebra generated by any one element is associative

Levels of Associativity

An algebra is:

- ***power-associative*** if the subalgebra generated by any one element is associative
- ***alternative*** if the subalgebra generated by any two elements is associative

Levels of Associativity

An algebra is:

- ***power-associative*** if the subalgebra generated by any one element is associative
- ***alternative*** if the subalgebra generated by any two elements is associative
- ***associative*** if the subalgebra generated by any three elements is associative

Commutator and Associator

- **Commutator** is an alternating bilinear map $[\cdot, \cdot] : \mathcal{A}^2 \rightarrow \mathcal{A}$ given by $[a, b] = ab - ba$. It switches sign whenever the two arguments are exchanged $[a, b] = -[b, a]$ or equivalently, that it vanishes when they are equal $[a, a] = 0$.

Commutator and Associator

- **Commutator** is an alternating bilinear map $[\cdot, \cdot] : \mathcal{A}^2 \rightarrow \mathcal{A}$ given by $[a, b] = ab - ba$. It switches sign whenever the two arguments are exchanged $[a, b] = -[b, a]$ or equivalently, that it vanishes when they are equal $[a, a] = 0$.
- **Associator** is a trilinear map $[\cdot, \cdot, \cdot] : \mathcal{A}^3 \rightarrow \mathcal{A}$ given by $[a, b, c] = (ab)c - a(bc)$.

Commutator and Associator

- **Commutator** is an alternating bilinear map $[\cdot, \cdot] : \mathcal{A}^2 \rightarrow \mathcal{A}$ given by $[a, b] = ab - ba$. It switches sign whenever the two arguments are exchanged $[a, b] = -[b, a]$ or equivalently, that it vanishes when they are equal $[a, a] = 0$.
- **Associator** is a trilinear map $[\cdot, \cdot, \cdot] : \mathcal{A}^3 \rightarrow \mathcal{A}$ given by $[a, b, c] = (ab)c - a(bc)$.
- The associator measures the failure of associativity just as the commutator measures the failure of commutativity.

Alternative Algebra

Theorem:

An algebra \mathcal{A} is alternative if and only if for all $a, b \in \mathcal{A}$ we have

$$(aa)b = a(ab), (ab)a = a(ba), (ba)a = b(aa)$$

[*Emil Artin, 1930s*]

Proposition:

Associator is alternating precisely when \mathcal{A} is alternative.

Basic Theorems

Theorem 1: \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} are the only normed division algebras. [*Hurwitz, 1898.*]

Basic Theorems

Theorem 1: \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} are the only normed division algebras. [*Hurwitz, 1898.*]

Theorem 2: \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} are the only alternative division algebras. [*Zorn, 1930.*]

Basic Theorems

Theorem 1: \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} are the only normed division algebras. [*Hurwitz, 1898.*]

Theorem 2: \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} are the only alternative division algebras. [*Zorn, 1930.*]

Theorem 3: All division algebras have dimension 1, 2, 4, or 8. [*Kervaire and Bott-Milnor, 1958.*]

Constructing the Octonions

- The octonions are an 8-dimensional algebra with basis $1, e_1, e_2, e_3, e_4, e_5, e_6, e_7$.
- The most elementary way to construct the octonions is to give their multiplication table.

Octonion Multiplication Table

	e_1	e_2	e_3	e_4	e_5	e_6	e_7
e_1	-1	e_4	e_7	$-e_2$	e_6	$-e_5$	$-e_3$
e_2	$-e_4$	-1	e_5	e_1	$-e_3$	e_7	$-e_6$
e_3	$-e_7$	$-e_5$	-1	e_6	e_2	$-e_4$	e_1
e_4	e_2	$-e_1$	$-e_6$	-1	e_7	e_3	$-e_5$
e_5	$-e_6$	e_3	$-e_2$	$-e_7$	-1	e_1	e_4
e_6	e_5	$-e_7$	e_4	e_3	$-e_1$	-1	e_2
e_7	e_3	e_6	$-e_1$	e_5	$-e_4$	$-e_2$	-1

Properties

- e_1, \dots, e_7 are square roots of -1

Properties

- e_1, \dots, e_7 are square roots of -1
- e_i and e_j anticommute when $i \neq j$: $e_i e_j = -e_j e_i$

Properties

- e_1, \dots, e_7 are square roots of -1
- e_i and e_j anticommute when $i \neq j$: $e_i e_j = -e_j e_i$
- the **index cycling** identity holds:
 $e_i e_j = e_k \Rightarrow e_{i+1} e_{j+1} = e_{k+1}$ (where we think of indices as living in \mathbb{Z}_7)

Properties

- e_1, \dots, e_7 are square roots of -1
- e_i and e_j anticommute when $i \neq j$: $e_i e_j = -e_j e_i$
- the **index cycling** identity holds:
 $e_i e_j = e_k \Rightarrow e_{i+1} e_{j+1} = e_{k+1}$ (where we think of indices as living in \mathbb{Z}_7)
- the **index doubling** identity holds:
 $e_i e_j = e_k \Rightarrow e_{2i} e_{2j} = e_{2k}$

Properties

- e_1, \dots, e_7 are square roots of -1
- e_i and e_j anticommute when $i \neq j$: $e_i e_j = -e_j e_i$
- the **index cycling** identity holds:
 $e_i e_j = e_k \Rightarrow e_{i+1} e_{j+1} = e_{k+1}$ (where we think of indices as living in \mathbb{Z}_7)
- the **index doubling** identity holds:
 $e_i e_j = e_k \Rightarrow e_{2i} e_{2j} = e_{2k}$
- nontrivial product: $e_1 e_2 = e_4$

Quaternions

The quaternions, \mathbb{H} , are a 4-dimensional algebra with basis $1, i, j, k$.

Multiplication properties:

Quaternions

The quaternions, \mathbb{H} , are a 4-dimensional algebra with basis $1, i, j, k$.

Multiplication properties:

- 1 is the multiplicative identity

Quaternions

The quaternions, \mathbb{H} , are a 4-dimensional algebra with basis $1, i, j, k$.

Multiplication properties:

- 1 is the multiplicative identity
- i, j and k are square roots of -1


Quaternions

The quaternions, \mathbb{H} , are a 4-dimensional algebra with basis $1, i, j, k$.


Multiplication properties:

- 1 is the multiplicative identity
- i, j and k are square roots of -1
- we have $ij = k$, $ji = -k$, and all identities obtained from these by cyclic permutations of (i, j, k)

Multiplication of Quaternions




Multiplication of Quaternions




Multiplication of Quaternions

$$ij = k$$




Multiplication of Quaternions

counterclockwise




Multiplication of Quaternions

$$ji = -k$$



How to multiply octonions?



Fano plane

Properties

- 1 is the multiplicative identity

Properties

- 1 is the multiplicative identity
- e_1, \dots, e_7 are square roots of -1

Properties


- 1 is the multiplicative identity
- e_1, \dots, e_7 are square roots of -1
- If e_i, e_j , and e_k are cyclically ordered in this way then $e_i e_j = e_k$, $e_j e_i = -e_k$.

Properties

- 1 is the multiplicative identity
- e_1, \dots, e_7 are square roots of -1
- If e_i, e_j , and e_k are cyclically ordered in this way then $e_i e_j = e_k$, $e_j e_i = -e_k$.
- the ***index doubling*** corresponds to rotating the picture a third of a turn


The Fano plane

- The Fano plane consists of 7 points and 7 lines.
- The 'lines' are the sides of the triangle, its altitudes, and the circle containing all the midpoints of the sides.
- It completely describes the algebra structure of the octonions.



Octonions \mathbb{O}

The Fano plane is the projective plane over the 2-element field \mathbb{Z}_2 .



Subalgebras of \mathbb{O}

- $1 \in \mathbb{O}$ - the origin in \mathbb{Z}_2^3 .

Subalgebras of \mathbb{O}

- $1 \in \mathbb{O}$ - the origin in \mathbb{Z}_2^3 .
- Planes through the origin give subalgebras isomorphic to the quaternions.

Subalgebras of \mathbb{O}

- $1 \in \mathbb{O}$ - the origin in \mathbb{Z}_2^3 .
- Planes through the origin give subalgebras isomorphic to the quaternions.
- Lines through the origin give subalgebras isomorphic to the complex numbers.

Subalgebras of \mathbb{O}

- $1 \in \mathbb{O}$ - the origin in \mathbb{Z}_2^3 .
- Planes through the origin give subalgebras isomorphic to the quaternions.
- Lines through the origin give subalgebras isomorphic to the complex numbers.
- The origin itself gives a subalgebra isomorphic to the real numbers.

The Cayley-Dickson Construction

The Cayley-Dickson construction gives an infinite sequence of algebras, doubling in dimension each time, with the normed division algebras \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} as the first four. Also, it explains why each one fits neatly inside the next.

Complex numbers

- $z = a + ib \in \mathbb{C} \iff (a, b) \in \mathbb{R}^2$

Complex numbers

- $z = a + ib \in \mathbb{C} \iff (a, b) \in \mathbb{R}^2$

- multiplication:

$$(a, b)(c, d) = (ac - db, ad + cb)$$

Complex numbers

- $z = a + ib \in \mathbb{C} \iff (a, b) \in \mathbb{R}^2$

- multiplication:

$$(a, b)(c, d) = (ac - db, ad + cb)$$

- conjugate:

$$(a, b)^* = (a, -b)$$

Quaternions

- $q \in \mathbb{H} \iff (a, b) \in \mathbb{C}^2$

Quaternions

- $q \in \mathbb{H} \iff (a, b) \in \mathbb{C}^2$

- multiplication:

$$(a, b)(c, d) = (ac - db^*, a^*d + cb)$$

Quaternions

- $q \in \mathbb{H} \iff (a, b) \in \mathbb{C}^2$

- multiplication:

$$(a, b)(c, d) = (ac - db^*, a^*d + cb)$$

- conjugate:

$$(a, b)^* = (a^*, -b)$$

Octonions

- We can define an octonion to be a pair of quaternions.

Octonions

- We can define an octonion to be a pair of quaternions.
- Multiplication and conjugate are defined in the same manner as for the quaternions.

Octonions

- We can define an octonion to be a pair of quaternions.
- Multiplication and conjugate are defined in the same manner as for the quaternions.
- Why isn't there an *infinite* sequence of division algebras, each one obtained from the preceding one by the Cayley-Dickson construction?

*-algebra

- An algebra \mathcal{A} equipped with a conjugation, that is, a real-linear map $*$: $\mathcal{A} \rightarrow \mathcal{A}$ with $a^{**} = a$, $(ab)^* = b^*a^*$ for all $a, b \in \mathcal{A}$ is called ***-algebra**.
- *-algebra \mathcal{A} is:
 - **real** if $a = a^*$ for every element $a \in \mathcal{A}$.
 - **nicely normed** if $a + a^* \in \mathbb{R}$ and $aa^* = a^*a > 0$ for all nonzero $a \in \mathcal{A}$.
- If \mathcal{A} is nicely normed and alternative, \mathcal{A} is a normed division algebra.

The General Construction

- Starting from any $*$ -algebra \mathcal{A} , the Cayley-Dickson construction gives a new $*$ -algebra \mathcal{A}' .
- Elements of \mathcal{A}' are pairs $(a, b) \in \mathcal{A}^2$.
- Multiplication:

$$(a, b)(c, d) = (ac - db^*, a^*d + cb)$$

- Conjugate:

$$(a, b)^* = (a^*, -b)$$

Propositions

P1: \mathcal{A}' is never real.

Propositions

P1: \mathcal{A}' is never real.

P2: \mathcal{A} is real (and thus commutative) \iff
 \mathcal{A}' is commutative.

Propositions

P1: \mathcal{A}' is never real.

P2: \mathcal{A} is real (and thus commutative) \iff
 \mathcal{A}' is commutative.

P3: \mathcal{A} is commutative and associative \iff
 \mathcal{A}' is associative.

Propositions

P1: \mathcal{A}' is never real.

P2: \mathcal{A} is real (and thus commutative) \iff
 \mathcal{A}' is commutative.

P3: \mathcal{A} is commutative and associative \iff
 \mathcal{A}' is associative.

P4: \mathcal{A} is associative and nicely normed \iff
 \mathcal{A}' is alternative and nicely normed.

Propositions

P1: \mathcal{A}' is never real.

P2: \mathcal{A} is real (and thus commutative) \iff
 \mathcal{A}' is commutative.

P3: \mathcal{A} is commutative and associative \iff
 \mathcal{A}' is associative.

P4: \mathcal{A} is associative and nicely normed \iff
 \mathcal{A}' is alternative and nicely normed.

P5: \mathcal{A} is nicely normed \iff \mathcal{A}' is nicely
normed.

Effects

\mathbb{R} is a real commutative associative nicely
normed $*$ -algebra



Effects

\mathbb{R} is a real commutative associative nicely
normed $*$ -algebra



\mathbb{C} is a commutative associative nicely normed
 $*$ -algebra



Effects

\mathbb{R} is a real commutative associative nicely
normed $*$ -algebra



\mathbb{C} is a commutative associative nicely normed
 $*$ -algebra



\mathbb{H} is an associative nicely normed $*$ -algebra



Effects

\mathbb{R} is a real commutative associative nicely
normed $*$ -algebra



\mathbb{C} is a commutative associative nicely normed
 $*$ -algebra



\mathbb{H} is an associative nicely normed $*$ -algebra




\mathbb{O} is an alternative nicely normed $*$ -algebra

Axioms of Projective Plane

- For any two distinct points, there is a unique line on which they both lie.
- For any two distinct lines, there is a unique point which lies on both of them.
- There exist four points, no three of which lie on the same line.
- There exist four lines, no three of which have the same point lying on them.



Examples of Projective Plane

Fano plane



Examples of Projective Plane

Fano plane



the real projective plane \mathbb{RP}^2

Points and Lines in $\mathbb{R}P^2$

- the points are lines through the origin in \mathbb{R}^3

Points and Lines in $\mathbb{R}P^2$

- the points are lines through the origin in \mathbb{R}^3
- each point $(x, y) \in \mathbb{R}^2$ determines a point in $\mathbb{R}P^2$, namely the line in \mathbb{R}^3 containing the origin and the point $(x, y, -1)$

Points and Lines in $\mathbb{R}P^2$

- the points are lines through the origin in \mathbb{R}^3
- each point $(x, y) \in \mathbb{R}^2$ determines a point in $\mathbb{R}P^2$, namely the line in \mathbb{R}^3 containing the origin and the point $(x, y, -1)$
- the ‘points at infinity’ are lines through the origin in \mathbb{R}^3 that do not intersect the plane $z = -1$

Points and Lines in $\mathbb{R}P^2$

- the points are lines through the origin in \mathbb{R}^3
- each point $(x, y) \in \mathbb{R}^2$ determines a point in $\mathbb{R}P^2$, namely the line in \mathbb{R}^3 containing the origin and the point $(x, y, -1)$
- the ‘points at infinity’ are lines through the origin in \mathbb{R}^3 that do not intersect the plane $z = -1$
- the lines are planes through the origin in \mathbb{R}^3

Axioms of Projective Space


- For any two distinct points p, q , there is a unique line pq on which they both lie.
- For any line, there are at least three points lying on this line.
- If a, b, c, d are distinct points and there is a point lying on both ab and cd , then there is a point lying on both ac and bd .

n -dimensional Projective Space

- \mathbb{K} - **skew field**: a ring such that every nonzero element has a left and right multiplicative inverse
- line through the origin: $L = \{\alpha x : \alpha \in \mathbb{K}\}$, where $x \in \mathbb{K}^{n+1}$ is nonzero
- plane through the origin: $P = \{\alpha x + \beta y : \alpha, \beta \in \mathbb{K}\}$, where $x, y \in \mathbb{K}^{n+1}$ are elements such that $\alpha x + \beta y = 0$ implies $\alpha, \beta = 0$
- projective n -space is of the form $\mathbb{K}P^n$ for some skew field \mathbb{K} only if $n > 2$


Axial Perspectivity

Axial perspectivity is the condition satisfied if and only if the points of intersection $L \cap L'$, $N \cap N'$ and $M \cap M'$, are collinear, on a line called *the axis of perspectivity*.




Central Perspectivity

Central perspectivity is the condition satisfied if and only if the three lines xx' , yy' and zz' are concurrent, at a point called ***the center of perspectivity***.



The Axiom of Desargues

In a projective space, two triangles are in perspective axially if and only if they are in perspective centrally.



A projective plane satisfying this axiom is called
Desarguesian.

Desarguesian Projective Planes and Skew Fields

We start with a projective plane P and try to reconstruct a skew field from it:

Desarguesian Projective Planes and Skew Fields

We start with a projective plane P and try to reconstruct a skew field from it:

- choose any line L and three distinct points on it - call them $0, 1$ and ∞

Desarguesian Projective Planes and Skew Fields

We start with a projective plane P and try to reconstruct a skew field from it:

- choose any line L and three distinct points on it - call them $0, 1$ and ∞
- set $\mathbb{K} = L - \{\infty\}$

Desarguesian Projective Planes and Skew Fields

We start with a projective plane P and try to reconstruct a skew field from it:

- choose any line L and three distinct points on it - call them $0, 1$ and ∞
- set $\mathbb{K} = L - \{\infty\}$
- define addition and multiplication of points in \mathbb{K}

Desarguesian Projective Planes and Skew Fields

We start with a projective plane P and try to reconstruct a skew field from it:

- choose any line L and three distinct points on it - call them $0, 1$ and ∞
- set $\mathbb{K} = L - \{\infty\}$
- define addition and multiplication of points in \mathbb{K}

The resulting structure $(\mathbb{K}, +, 0, \cdot, 1)$ will not be a skew field and it will depend in a nontrivial way on the choices made,

Desarguesian Projective Planes and Skew Fields

We start with a projective plane P and try to reconstruct a skew field from it:

- choose any line L and three distinct points on it - call them $0, 1$ and ∞
- set $\mathbb{K} = L - \{\infty\}$
- define addition and multiplication of points in \mathbb{K}

The resulting structure $(\mathbb{K}, +, 0, \cdot, 1)$ will not be a skew field and it will depend in a nontrivial way on the choices made, but if we assume the axiom of Desargues, these problems go away.

Brief History

- Projective geometry was very fashionable in the 1800s, with such worthies as *Poncelet*, *Brianchon*, *Steiner* and *von Staudt* making important contributions.

Brief History

- Projective geometry was very fashionable in the 1800s, with such worthies as *Poncelet*, *Brianchon*, *Steiner* and *von Staudt* making important contributions.
- In 1933 *Ruth Moufang* constructed a remarkable example of a non-Desarguesian projective plane using the octonions.

Brief History

- Projective geometry was very fashionable in the 1800s, with such worthies as *Poncelet*, *Brianchon*, *Steiner* and *von Staudt* making important contributions.
- In 1933 *Ruth Moufang* constructed a remarkable example of a non-Desarguesian projective plane using the octonions.
- The 1930s also saw the rise of another reason for interest in projective geometry: quantum mechanics!
However, in quantum mechanics, observables do not form an associative algebra.

Brief History

- Projective geometry was very fashionable in the 1800s, with such worthies as *Poncelet*, *Brianchon*, *Steiner* and *von Staudt* making important contributions.
- In 1933 *Ruth Moufang* constructed a remarkable example of a non-Desarguesian projective plane using the octonions.
- The 1930s also saw the rise of another reason for interest in projective geometry: quantum mechanics!
However, in quantum mechanics, observables do not form an associative algebra.
- In 1932, *Pascual Jordan* attempted to understand this situation better by isolating the bare minimum axioms that an 'algebra of observables' should satisfy.

Brief History

- Projective geometry was very fashionable in the 1800s, with such worthies as *Poncelet*, *Brianchon*, *Steiner* and *von Staudt* making important contributions.
- In 1933 *Ruth Moufang* constructed a remarkable example of a non-Desarguesian projective plane using the octonions.
- The 1930s also saw the rise of another reason for interest in projective geometry: quantum mechanics!
However, in quantum mechanics, observables do not form an associative algebra.
- In 1932, *Pascual Jordan* attempted to understand this situation better by isolating the bare minimum axioms that an ‘algebra of observables’ should satisfy.
- In 1934, *Jordan* published a paper with *von Neumann* and *Wigner* classifying all formally real Jordan algebras

Formally Real Jordan Algebra

- *A formally real Jordan algebra* is a commutative and power-associative algebra satisfying

$$a_1^2 + \cdots + a_n^2 = 0 \quad \implies \quad a_1 = \cdots = a_n = 0$$

for all n .

Formally Real Jordan Algebra

- **A formally real Jordan algebra** is a commutative and power-associative algebra satisfying

$$a_1^2 + \cdots + a_n^2 = 0 \quad \implies \quad a_1 = \cdots = a_n = 0$$

for all n .

- The last condition gives the algebra a partial ordering: if we write $a \leq b$ when the element $b - a$ is a sum of squares, it says that $a \leq b$ and $b \leq a$ imply $a = b$.

Jordan Algebra

- One can construct a commutative but nonassociative product:

$$a \circ b = \frac{1}{2}((a + b)^2 - a^2 - b^2) = \frac{1}{2}(ab + ba).$$

Jordan Algebra

- One can construct a commutative but nonassociative product:

$$a \circ b = \frac{1}{2}((a + b)^2 - a^2 - b^2) = \frac{1}{2}(ab + ba).$$

- Any formally real Jordan algebra satisfies the identity $a \circ (b \circ a^2) = (a \circ b) \circ a^2$ for all elements a and b .

Jordan Algebra

- One can construct a commutative but nonassociative product:

$$a \circ b = \frac{1}{2}((a + b)^2 - a^2 - b^2) = \frac{1}{2}(ab + ba).$$

- Any formally real Jordan algebra satisfies the identity $a \circ (b \circ a^2) = (a \circ b) \circ a^2$ for all elements a and b .
- Any commutative algebra satisfying this identity is called a **Jordan algebra**.
Jordan algebras are automatically power-associative.

Classification

The simple formally real Jordan algebras consist of 4 infinite families and one exception:

- The algebra $\mathfrak{h}_n(\mathbb{R})$ with the product $a \circ b = \frac{1}{2}(ab + ba)$.
- The algebra $\mathfrak{h}_n(\mathbb{C})$ with the product $a \circ b = \frac{1}{2}(ab + ba)$.
- The algebra $\mathfrak{h}_n(\mathbb{H})$ with the product $a \circ b = \frac{1}{2}(ab + ba)$.
- The algebra $\mathbb{R}^n \oplus \mathbb{R}$ with the product
 $(v, \alpha) \circ (w, \beta) = (\alpha w + \beta v, \langle v, w \rangle + \alpha\beta)$.
- The algebra $\mathfrak{h}_3(\mathbb{O})$ with the product $a \circ b = \frac{1}{2}(ab + ba)$.

Relation between $\mathfrak{h}_n(\mathbb{C})$ and $\mathbb{C}P^n$

- projections in $\mathfrak{h}_n(\mathbb{C})$ correspond to subspaces of \mathbb{C}^n
- the projections onto 1-dimensional subspaces correspond to points in $\mathbb{C}P^n$
- the projections onto 2-dimensional subspaces correspond to lines in $\mathbb{C}P^n$

Constructing Projective Space

We can then try to construct a projective space whose points are the rank-1 projections and whose lines are the rank-2 projections, with the relation of 'lying on' given by the partial order \leq .

Constructing Projective Space

We can then try to construct a projective space whose points are the rank-1 projections and whose lines are the rank-2 projections, with the relation of ‘lying on’ given by the partial order \leq .

- from $\mathfrak{h}_n(\mathbb{R})$, $\mathfrak{h}_n(\mathbb{C})$ or $\mathfrak{h}_n(\mathbb{H})$, we obtain the projective spaces $\mathbb{R}P^n$, $\mathbb{C}P^n$ and $\mathbb{H}P^n$, respectively for $n \geq 2$.

Constructing Projective Space

We can then try to construct a projective space whose points are the rank-1 projections and whose lines are the rank-2 projections, with the relation of 'lying on' given by the partial order \leq .

- from $\mathfrak{h}_n(\mathbb{R})$, $\mathfrak{h}_n(\mathbb{C})$ or $\mathfrak{h}_n(\mathbb{H})$, we obtain the projective spaces $\mathbb{R}P^n$, $\mathbb{C}P^n$ and $\mathbb{H}P^n$, respectively for $n \geq 2$.
- from *the spin factor* $\mathbb{R}^n \oplus \mathbb{R}$, for $n \geq 2$, we obtain a series of 1-dimensional projective spaces related to Lorentzian geometry.

Constructing Projective Space

We can then try to construct a projective space whose points are the rank-1 projections and whose lines are the rank-2 projections, with the relation of 'lying on' given by the partial order \leq .

- from $\mathfrak{h}_n(\mathbb{R})$, $\mathfrak{h}_n(\mathbb{C})$ or $\mathfrak{h}_n(\mathbb{H})$, we obtain the projective spaces $\mathbb{R}P^n$, $\mathbb{C}P^n$ and $\mathbb{H}P^n$, respectively for $n \geq 2$.
- from *the spin factor* $\mathbb{R}^n \oplus \mathbb{R}$, for $n \geq 2$, we obtain a series of 1-dimensional projective spaces related to Lorentzian geometry.
- starting with *the exceptional Jordan algebra* $\mathfrak{h}_3(\mathbb{O})$, we get the projective plane discovered by Moufang $\mathbb{O}P^2$.

Projective Lines

- A one-dimensional projective space is called a *projective line*.

Projective Lines

- A one-dimensional projective space is called a *projective line*.
- Since octonions are not associative, the problem of defining the projective line over \mathbb{O} arises.

Projective Lines

- A one-dimensional projective space is called a *projective line*.
- Since octonions are not associative, the problem of defining the projective line over \mathbb{O} arises.
- A square matrix with entries in the $*$ -algebra \mathcal{A} is *hermitian* if it equals its conjugate transpose.

Projective Lines

- A one-dimensional projective space is called a *projective line*.
- Since octonions are not associative, the problem of defining the projective line over \mathbb{O} arises.
- A square matrix with entries in the $*$ -algebra \mathcal{A} is *hermitian* if it equals its conjugate transpose.
- Let $\mathfrak{h}_n(\mathcal{A})$ stand for the hermitian $n \times n$ matrices with entries in \mathcal{A} .

Projections in $\mathfrak{h}_2(\mathbb{O})$

- Besides the trivial projections 0 and 1, they are all of the form:

$$\begin{pmatrix} x^* \\ y^* \end{pmatrix} \begin{pmatrix} x & y \end{pmatrix} = \begin{pmatrix} x^*x & x^*y \\ y^*x & y^*y \end{pmatrix}$$

where $(x, y) \in \mathbb{O}^2$ has $\|x\|^2 + \|y\|^2 = 1$.

- These nontrivial projections all have rank 1, so they are the points of one line
- It is easy to check that the axioms for a projective space hold.

- We have succeeded in creating the *projective line over \mathbb{O}* and that is $\mathbb{O}P^1$.

- We have succeeded in creating the **projective line over \mathbb{O}** and that is $\mathbb{O}P^1$.
- Given any nonzero element $(x, y) \in \mathbb{O}^2$, we can normalize it and then get a point in $\mathbb{O}P^1$, which we call $[(x, y)]$.

- We have succeeded in creating the **projective line over \mathbb{O}** and that is $\mathbb{O}P^1$.
- Given any nonzero element $(x, y) \in \mathbb{O}^2$, we can normalize it and then get a point in $\mathbb{O}P^1$, which we call $[(x, y)]$.
- Define an equivalence relation on nonzero elements of \mathbb{O}^2 by $(x, y) \sim (x', y') \iff [(x, y)] = [(x', y')]$.

- We have succeeded in creating the **projective line over \mathbb{O}** and that is $\mathbb{O}P^1$.
- Given any nonzero element $(x, y) \in \mathbb{O}^2$, we can normalize it and then get a point in $\mathbb{O}P^1$, which we call $[(x, y)]$.
- Define an equivalence relation on nonzero elements of \mathbb{O}^2 by $(x, y) \sim (x', y') \iff [(x, y)] = [(x', y')]$.
- We call an equivalence class for this relation a **line through the origin** in \mathbb{O}^2 .

Lines through the Origin

- We can then identify points in $\mathbb{O}P^1$ with lines through the origin in \mathbb{O}^2 .

Lines through the Origin

- We can then identify points in $\mathbb{O}P^1$ with lines through the origin in \mathbb{O}^2 .
- $\{(\alpha x, \alpha y) : \alpha \in \mathbb{O}\}$ is **not** line through the origin containing (x, y) !!!

Lines through the Origin

- We can then identify points in $\mathbb{O}P^1$ with lines through the origin in \mathbb{O}^2 .
- $\{(\alpha x, \alpha y) : \alpha \in \mathbb{O}\}$ is **not** line through the origin containing (x, y) !!!
- $(x, y) \sim (y^{-1}x, 1)$ when $y \neq 0$ and $(x, y) \sim (1, x^{-1}y)$ when $x \neq 0$, so the corresponding lines are $\{(\alpha, \alpha(x^{-1}y)) : \alpha \in \mathbb{O}\}$ and $\{(\alpha(y^{-1}x), \alpha) : \alpha \in \mathbb{O}\}$

Lines through the Origin

- We can then identify points in $\mathbb{O}P^1$ with lines through the origin in \mathbb{O}^2 .
- $\{(\alpha x, \alpha y) : \alpha \in \mathbb{O}\}$ is **not** line through the origin containing (x, y) !!!
- $(x, y) \sim (y^{-1}x, 1)$ when $y \neq 0$ and $(x, y) \sim (1, x^{-1}y)$ when $x \neq 0$, so the corresponding lines are $\{(\alpha, \alpha(x^{-1}y)) : \alpha \in \mathbb{O}\}$ and $\{(\alpha(y^{-1}x), \alpha) : \alpha \in \mathbb{O}\}$
- In particular, the line through the origin containing (x, y) is always a real vector space isomorphic to \mathbb{O} .

Manifolds

- Let \mathbb{K} denote a normed division algebra.

Manifolds

- Let \mathbb{K} denote a normed division algebra.
- We can cover $\mathbb{K}P^1$ with two coordinate charts: one containing all points of the form $[(x, 1)]$, the other containing all points of the form $[(1, y)]$.

Manifolds

- Let \mathbb{K} denote a normed division algebra.
- We can cover $\mathbb{K}P^1$ with two coordinate charts: one containing all points of the form $[(x, 1)]$, the other containing all points of the form $[(1, y)]$.
- It is easy to check that $[(x, 1)] = [(1, y)] \Leftrightarrow y = x^{-1}$, so the transition function from the first chart to the second is the map $x \mapsto x^{-1}$.

Manifolds


- Let \mathbb{K} denote a normed division algebra.
- We can cover $\mathbb{K}P^1$ with two coordinate charts: one containing all points of the form $[(x, 1)]$, the other containing all points of the form $[(1, y)]$.
- It is easy to check that $[(x, 1)] = [(1, y)] \Leftrightarrow y = x^{-1}$, so the transition function from the first chart to the second is the map $x \mapsto x^{-1}$.
- Since this transition function and its inverse are smooth on the intersection of the two charts, $\mathbb{K}P^1$ becomes a smooth manifold.

Complex Case

- $\mathbb{C}P^1$ is just the *Riemann sphere*.

Complex Case

- $\mathbb{C}P^1$ is just the *Riemann sphere*.
- the map $x \mapsto [(x, 1)]$ is given by stereographic projection:



where we choose the sphere to have diameter 1.

Conformal Inversion

- This map from \mathbb{C} to $\mathbb{C}P^1$ is one-to-one and almost onto, missing only the point at infinity, or *north pole*.

Conformal Inversion

- This map from \mathbb{C} to $\mathbb{C}P^1$ is one-to-one and almost onto, missing only the point at infinity, or *north pole*.
- Similarly, the map $y \mapsto [(1, y)]$ misses only the *south pole*.

Conformal Inversion

- This map from \mathbb{C} to $\mathbb{C}P^1$ is one-to-one and almost onto, missing only the point at infinity, or *north pole*.
- Similarly, the map $y \mapsto [(1, y)]$ misses only the *south pole*.
- Composing the first map with the inverse of the second, we get the map $x \mapsto x^{-1}$, which goes by the name of *conformal inversion*.

Riemann Sphere

- The southern hemisphere of the Riemann sphere consists of all points $[(x, 1)]$ with $\|x\| \leq 1$, while the northern hemisphere consists of all $[(1, y)]$ with $\|y\| \leq 1$.
- Unit complex numbers x give points $[(x, 1)] = [(1, x^{-1})]$ on the equator.

Generalizing the Idea

Smooth manifold $\mathbb{K}P^1$ is just a sphere with dimension equal to that of \mathbb{K} :

$$\mathbb{R}P^1 \cong S^1$$

$$\mathbb{C}P^1 \cong S^2$$

$$\mathbb{H}P^1 \cong S^4$$

$$\mathbb{O}P^1 \cong S^8$$

Sphere Description

- We can think of it as the one-point compactification of \mathbb{K} .

Sphere Description

- We can think of it as the one-point compactification of \mathbb{K} .
- The *southern hemisphere*, *northern hemisphere* and *equator* of \mathbb{K} have descriptions exactly like those given above for the complex case.

Sphere Description

- We can think of it as the one-point compactification of \mathbb{K} .
- The *southern hemisphere*, *northern hemisphere* and *equator* of \mathbb{K} have descriptions exactly like those given above for the complex case.
- Also, as in the complex case, the maps $x \mapsto [(x, 1)]$ and $y \mapsto [(1, y)]$ are angle-preserving with respect to the usual Euclidean metric on \mathbb{K} and the round metric on the sphere.

Line Bundles

- $\mathbb{K}P^1$ is equipped with a vector bundle whose fiber over the point $[(x, y)]$ is the line through the origin corresponding to this point.

Line Bundles

- $\mathbb{K}P^1$ is equipped with a vector bundle whose fiber over the point $[(x, y)]$ is the line through the origin corresponding to this point.
- This bundle is called *the canonical line bundle*, $L_{\mathbb{K}}$.

Line Bundles

- $\mathbb{K}P^1$ is equipped with a vector bundle whose fiber over the point $[(x, y)]$ is the line through the origin corresponding to this point.
- This bundle is called *the canonical line bundle*, $L_{\mathbb{K}}$.
- When we are working with a particular division algebra, ‘line’ means a copy of this division algebra, so if we think of them as real vector bundles, $L_{\mathbb{R}}$, $L_{\mathbb{C}}$, $L_{\mathbb{H}}$ and $L_{\mathbb{O}}$ have dimensions 1, 2, 4 and 8, respectively.

Building Canonical Line Bundles

- Any k -dimensional real vector bundle over S^n can be formed by taking trivial bundles over the northern and southern hemispheres and gluing them together along the equator via a map $f : S^{n-1} \rightarrow \mathbf{O}(n)$

Building Canonical Line Bundles

- Any k -dimensional real vector bundle over S^n can be formed by taking trivial bundles over the northern and southern hemispheres and gluing them together along the equator via a map $f : S^{n-1} \rightarrow \mathbf{O}(n)$
- We must therefore be able to build the canonical line bundles $L_{\mathbb{R}}, L_{\mathbb{C}}, L_{\mathbb{H}}$ and $L_{\mathbb{O}}$ using maps:

$$f_{\mathbb{R}} : S^0 \rightarrow \mathbf{O}(1)$$

$$f_{\mathbb{C}} : S^1 \rightarrow \mathbf{O}(2)$$

$$f_{\mathbb{H}} : S^3 \rightarrow \mathbf{O}(4)$$

$$f_{\mathbb{O}} : S^7 \rightarrow \mathbf{O}(8)$$

What Are These Maps?

- In the southern hemisphere of $\mathbb{K}P^1$, we can identify any fiber of $L_{\mathbb{K}}$ with \mathbb{K} by mapping the point $(\alpha x, \alpha)$ in the line $[(x, 1)]$ to the element $\alpha \in \mathbb{K}$.

What Are These Maps?

- In the southern hemisphere of $\mathbb{K}P^1$, we can identify any fiber of $L_{\mathbb{K}}$ with \mathbb{K} by mapping the point $(\alpha x, \alpha)$ in the line $[(x, 1)]$ to the element $\alpha \in \mathbb{K}$.
- Similarly, we can trivialize this bundle over the northern hemisphere by mapping the point $(\beta, \beta y)$ in the line $[(1, y)]$ to the element $\beta \in \mathbb{K}$.

What Are These Maps?

- In the southern hemisphere of $\mathbb{K}P^1$, we can identify any fiber of $L_{\mathbb{K}}$ with \mathbb{K} by mapping the point $(\alpha x, \alpha)$ in the line $[(x, 1)]$ to the element $\alpha \in \mathbb{K}$.
- Similarly, we can trivialize this bundle over the northern hemisphere by mapping the point $(\beta, \beta y)$ in the line $[(1, y)]$ to the element $\beta \in \mathbb{K}$.
- If $x \in \mathbb{K}$ has norm one, $[(x, 1)] = [(1, x^{-1})]$ is a point on the equator, so we get two trivializations of the fiber over this point: if $(\alpha x, \alpha) = (\beta, \beta x^{-1})$ then $\beta = \alpha x$.

What Are These Maps?

- In the southern hemisphere of $\mathbb{K}P^1$, we can identify any fiber of $L_{\mathbb{K}}$ with \mathbb{K} by mapping the point $(\alpha x, \alpha)$ in the line $[(x, 1)]$ to the element $\alpha \in \mathbb{K}$.
- Similarly, we can trivialize this bundle over the northern hemisphere by mapping the point $(\beta, \beta y)$ in the line $[(1, y)]$ to the element $\beta \in \mathbb{K}$.
- If $x \in \mathbb{K}$ has norm one, $[(x, 1)] = [(1, x^{-1})]$ is a point on the equator, so we get two trivializations of the fiber over this point: if $(\alpha x, \alpha) = (\beta, \beta x^{-1})$ then $\beta = \alpha x$.
- The map $\alpha \mapsto \beta$ is thus right multiplication by x .

The Importance of $f_{\mathbb{K}}$

- $f_{\mathbb{K}} : S^{n-1} \rightarrow \mathbf{O}(n)$ is just the map sending any norm-one element $x \in \mathbb{K}$ to the operation of right multiplication by x .

The Importance of $f_{\mathbb{K}}$

- $f_{\mathbb{K}} : S^{n-1} \rightarrow \mathbf{O}(n)$ is just the map sending any norm-one element $x \in \mathbb{K}$ to the operation of right multiplication by x .
- Using the obvious inclusions $\mathbf{O}(n) \hookrightarrow \mathbf{O}(n+1)$, we obtain a topological group called $\mathbf{O}(\infty)$

The Importance of $f_{\mathbb{K}}$

- $f_{\mathbb{K}} : S^{n-1} \rightarrow \mathbf{O}(n)$ is just the map sending any norm-one element $x \in \mathbb{K}$ to the operation of right multiplication by x .
- Using the obvious inclusions $\mathbf{O}(n) \hookrightarrow \mathbf{O}(n+1)$, we obtain a topological group called $\mathbf{O}(\infty)$
- Since $\mathbf{O}(n)$ is included in $\mathbf{O}(\infty)$, we can think of $f_{\mathbb{K}}$ as a map from S^{n-1} to $\mathbf{O}(\infty)$.

The Importance of $f_{\mathbb{K}}$

Homotopy class $[f_{\mathbb{K}}]$ has the following marvelous property, calculated by *Raoul Bott* in 1957:

$$[f_{\mathbb{R}}] \text{ generates } \pi_0(\mathbf{O}(\infty)) \cong \mathbb{Z}_2$$

$$[f_{\mathbb{C}}] \text{ generates } \pi_1(\mathbf{O}(\infty)) \cong \mathbb{Z}_2$$

$$[f_{\mathbb{H}}] \text{ generates } \pi_3(\mathbf{O}(\infty)) \cong \mathbb{Z}$$

$$[f_{\mathbb{O}}] \text{ generates } \pi_7(\mathbf{O}(\infty)) \cong \mathbb{Z}$$

Bott Periodicity

Bott proved that the homotopy groups of the topological group $\mathbf{O}(\infty)$ repeat with period 8:

$$\pi_{i+8}(\mathbf{O}(\infty)) \cong \pi_i(\mathbf{O}(\infty)).$$

He also computed the first 8:

$$\pi_0(\mathbf{O}(\infty)) \cong \mathbb{Z}_2$$

$$\pi_4(\mathbf{O}(\infty)) \cong 0$$

$$\pi_1(\mathbf{O}(\infty)) \cong \mathbb{Z}_2$$

$$\pi_5(\mathbf{O}(\infty)) \cong 0$$

$$\pi_2(\mathbf{O}(\infty)) \cong 0$$

$$\pi_6(\mathbf{O}(\infty)) \cong 0$$

$$\pi_3(\mathbf{O}(\infty)) \cong \mathbb{Z}$$

$$\pi_7(\mathbf{O}(\infty)) \cong \mathbb{Z}$$

References:

- "*On Octonions*" by John C. Baez,
Bull. Amer. Math. Soc. 39 (2002), 145-205.
- "*On Quaternions and Octonions: Their Geometry, Arithmetic, and Symmetry*" by John H. Conway and Derek A. Smith,
Bull. Amer. Math. Soc. 42 (2005), 229-243.
- "*Ubiquitous octonions*" by John C. Baez,
Plus Magazine, Issue 33, January 2005



Thank you for the attention!

Questions?