

Quaternions and their geometry

Eva Černohorská

Rapotín

27 December 2010

Overview

- ▶ Motivation
 - ▶ \mathbb{C} and 2D geometry
 - ▶ History
- ▶ Quaternions
 - ▶ Definition
 - ▶ \mathbb{H} and 3D geometry
- ▶ Generalization

Complex numbers

► $\mathbb{C} = \{z \mid z = x + iy, x, y \in \mathbb{R}, i^2 = -1\}$

Complex numbers

- ▶ $\mathbb{C} = \{z | z = x + iy, x, y \in \mathbb{R}, i^2 = -1\}$
- ▶ *Euclidean norm* $|z| = |x + iy| = \sqrt{x^2 + y^2}$

Complex numbers

- ▶ $\mathbb{C} = \{z \mid z = x + iy, x, y \in \mathbb{R}, i^2 = -1\}$
- ▶ *Euclidean norm* $|z| = |x + iy| = \sqrt{x^2 + y^2}$
 - ▶ $|z_1 z_2| = |z_1| |z_2|$

Complex numbers

- ▶ $\mathbb{C} = \{z \mid z = x + iy, x, y \in \mathbb{R}, i^2 = -1\}$
- ▶ *Euclidean norm* $|z| = |x + iy| = \sqrt{x^2 + y^2}$
 - ▶ $|z_1 z_2| = |z_1| |z_2|$
 - ▶ multiplying z by z_0 multiplies the norm $|z|$ by $\sqrt{|z_0|}$

Complex numbers

- ▶ $\mathbb{C} = \{z | z = x + iy, x, y \in \mathbb{R}, i^2 = -1\}$
- ▶ *Euclidean norm* $|z| = |x + iy| = \sqrt{x^2 + y^2}$
 - ▶ $|z_1 z_2| = |z_1| |z_2|$
 - ▶ multiplying z by z_0 multiply the norm $|z|$ by $\sqrt{|z_0|}$
 - ▶ when $|z_0| = 1$ (z_0 is a *complex unit*) the multiplication by z_0 is *isometry*

Rotation

- ▶ congruence which can be obtained by a continuous motion from the identity

Rotation

- ▶ congruence which can be obtained by a continuous motion from the identity
- ▶ standard formula

$$x' = x \cos \theta - y \sin \theta$$

$$y' = x \sin \theta + y \cos \theta$$

Rotation

- ▶ congruence which can be obtained by a continuous motion from the identity

- ▶ standard formula

$$x' = x \cos \theta - y \sin \theta$$

$$y' = x \sin \theta + y \cos \theta$$

- ▶ follows from multiplication of complex numbers

$$(\cos \theta + i \sin \theta)(x + iy) = x \cos \theta - y \sin \theta + i(x \sin \theta + y \cos \theta)$$


Rotation

- ▶ congruence which can be obtained by a continuous motion from the identity

- ▶ standard formula

$$x' = x \cos \theta - y \sin \theta$$

$$y' = x \sin \theta + y \cos \theta$$



- ▶ follows from multiplication of complex numbers

$$(\cos \theta + i \sin \theta)(x + iy) = x \cos \theta - y \sin \theta + i(x \sin \theta + y \cos \theta)$$

Reflection

- ▶ standard formula

$$x' = x$$

$$y' = -y$$

Reflection

- ▶ standard formula

$$x' = x$$

$$y' = -y$$


- ▶ corresponds to conjugation $\overline{x + iy} = x - iy$

Reflection

- ▶ standard formula

$$x' = x$$

$$y' = -y$$




- ▶ corresponds to conjugation $\overline{x + iy} = x - iy$

Reflection

- ▶ standard formula

$$x' = x$$

$$y' = -y$$



- ▶ corresponds to conjugation $\overline{x + iy} = x - iy$

Theorem

If u is a complex unit then the map $z \rightarrow zu$ is a rotation, while $z \rightarrow z\bar{u}$ is a reflection.

GO_2 - General orthogonal group

SO_2 - Special orthogonal group

GO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$

GO_2 - General orthogonal group

SO_2 - Special orthogonal group

GO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$
- ▶ the set of all isometries of Euclidean plane fixing the origin

GO_2 - General orthogonal group

SO_2 - Special orthogonal group

GO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$
- ▶ the set of all isometries of Euclidean plane fixing the origin
- ▶ the determinant of every matrix from GO_2 is ± 1

GO_2 - General orthogonal group

SO_2 - Special orthogonal group

GO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$
- ▶ the set of all isometries of Euclidean plane fixing the origin
- ▶ the determinant of every matrix from GO_2 is ± 1

SO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$ with determinant 1

GO_2 - General orthogonal group

SO_2 - Special orthogonal group

GO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$
- ▶ the set of all isometries of Euclidean plane fixing the origin
- ▶ the determinant of every matrix from GO_2 is ± 1

SO_2

- ▶ the set of 2×2 matrices A satisfying $AA^T = A^T A = I$ with determinant 1
- ▶ the set of all isometries of Euclidean plane fixing the origin preserving handedness

Theorem

Every element from GO_2 is a rotation or the reflection.

Theorem

Every element from GO_2 is a rotation or the reflection.

Proof.

Let $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ be the orthogonal matrix.

First orthogonality condition ($a_{11}^2 + a_{12}^2 = 1$) tells that $a_{11} = \cos \theta$ and $a_{12} = \sin \theta$ for some θ .

Second orthogonality condition ($a_{11}a_{21} + a_{12}a_{22} = 0$) is equivalent to $a_{21} = -a_{22} \tan \theta$ and

from the third condition ($a_{21}^2 + a_{22}^2 = 1$) follows that $a_{22} = \pm \cos \theta$ and $a_{21} = \mp \sin \theta$. □

GO_2, SO_2

rotation	reflection
$\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$	$\begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$
determinant $+1$	determinant -1
SO_2	$GO_2 \setminus SO_2$
$z \rightarrow zu$	$z \rightarrow z\bar{u}$

History

- ▶ 1748 Euler 4-squares identity

History

- ▶ 1748 Euler 4-squares identity
- ▶ 1840 Rodrigues parametrization of general rotation by four parameters

History

- ▶ 1748 Euler 4-squares identity
- ▶ 1840 Rodrigues parametrization of general rotation by four parameters
- ▶ 1835 Hamilton construction of quaternions as an algebra

William Rowan Hamilton 1805 - 1865



William Rowan Hamilton 1805 - 1865



- ▶ 1835 - representation of complex numbers as pair of real numbers, relation between \mathbb{C} and 2D geometry

William Rowan Hamilton 1805 - 1865



- ▶ 1835 - representation of complex numbers as pair of real numbers, relation between \mathbb{C} and 2D geometry
- ▶ tried to invent bigger algebra with the similar role in 3D geometry

William Rowan Hamilton 1805 - 1865



- ▶ 1835 - representation of complex numbers as pair of real numbers, relation between \mathbb{C} and 2D geometry
- ▶ tried to invent bigger algebra with the similar role in 3D geometry

"Every morning on my coming down to breakfast, your little brother William Edwin, and yourself, used to ask me: 'Well, Papa, can you multiply triplets?' Whereto I was always obliged to reply, with a sad shake of the head: 'No, I can only add and subtract them.'"

William Rowan Hamilton 1805 - 1865



- ▶ 1835 - representation of complex numbers as pair of real numbers, relation between \mathbb{C} and 2D geometry
- ▶ tried to invent bigger algebra with the similar role in 3D geometry

"Every morning on my coming down to breakfast, your little brother William Edwin, and yourself, used to ask me: 'Well, Papa, can you multiply triplets?' Whereto I was always obliged to reply, with a sad shake of the head: 'No, I can only add and subtract them.'"

- ▶ 16 October 1843 invention of quaternions

William Rowan Hamilton 1805 - 1865



- ▶ 1835 - representation of complex numbers as pair of real numbers, relation between \mathbb{C} and 2D geometry
- ▶ tried to invent bigger algebra with the similar role in 3D geometry

“Every morning on my coming down to breakfast, your little brother William Edwin, and yourself, used to ask me: ‘Well, Papa, can you multiply triplets?’ Whereto I was always obliged to reply, with a sad shake of the head: ‘No, I can only add and subtract them.’”

- ▶ 16 October 1843 invention of quaternions
- ▶ carved equation into the stone of the Brougham Bridge

William Rowan Hamilton 1805 - 1865




- ▶ 1835 - representation of complex numbers as pair of real numbers, relation between \mathbb{C} and 2D geometry
- ▶ tried to invent bigger algebra with the similar role in 3D geometry

“Every morning on my coming down to breakfast, your little brother William Edwin, and yourself, used to ask me: ‘Well, Papa, can you multiply triplets?’ Whereto I was always obliged to reply, with a sad shake of the head: ‘No, I can only add and subtract them.’”

- ▶ 16 October 1843 invention of quaternions
- ▶ carved equation into the stone of the Brougham Bridge
- ▶ rest of life with quaternions and their application to geometry

Quaternions - definition


$$\mathbb{H} = \{q | q = a+bi+cj+dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$


- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$,
 $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$


- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$, $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$
- ▶ multiplication:
 $[a, \mathbf{v}][a', \mathbf{v}'] = qq' = [aa' - \mathbf{v} \cdot \mathbf{v}', \mathbf{v} \times \mathbf{v}' + a\mathbf{v}' + a'\mathbf{v}]$



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$


- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$, $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$
- ▶ multiplication:
 $[a, \mathbf{v}][a', \mathbf{v}'] = qq' = [aa' - \mathbf{v} \cdot \mathbf{v}', \mathbf{v} \times \mathbf{v}' + a\mathbf{v}' + a'\mathbf{v}]$
- ▶ the cross product is not commutative, so the multiplication of quaternions is not



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$


- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$, $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$
- ▶ multiplication:
 $[a, \mathbf{v}][a', \mathbf{v}'] = qq' = [aa' - \mathbf{v} \cdot \mathbf{v}', \mathbf{v} \times \mathbf{v}' + a\mathbf{v}' + a'\mathbf{v}]$
- ▶ the cross product is not commutative, so the multiplication of quaternions is not
- ▶ the multiplication of quaternions is



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$


- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$, $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$
- ▶ multiplication:
 $[a, \mathbf{v}][a', \mathbf{v}'] = qq' = [aa' - \mathbf{v} \cdot \mathbf{v}', \mathbf{v} \times \mathbf{v}' + a\mathbf{v}' + a'\mathbf{v}]$
- ▶ the cross product is not commutative, so the multiplication of quaternions is not
- ▶ the multiplication of quaternions is
 - ▶ associative



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$


- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$, $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$
- ▶ multiplication:
 $[a, \mathbf{v}][a', \mathbf{v}'] = qq' = [aa' - \mathbf{v} \cdot \mathbf{v}', \mathbf{v} \times \mathbf{v}' + a\mathbf{v}' + a'\mathbf{v}]$
- ▶ the cross product is not commutative, so the multiplication of quaternions is not
- ▶ the multiplication of quaternions is
 - ▶ associative
 - ▶ distributive with respect to addition



Quaternions - definition

$$\mathbb{H} = \{q | q = a + bi + cj + dk, a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\}$$

- ▶ notation: $\mathbb{H} \ni q = a + bi + cj + dk = [a, \mathbf{v}]$, where $s \in \mathbb{R}$, $\mathbf{v} = (b, c, d) \in \mathbb{R}^3$
- ▶ multiplication:
 $[a, \mathbf{v}][a', \mathbf{v}'] = qq' = [aa' - \mathbf{v} \cdot \mathbf{v}', \mathbf{v} \times \mathbf{v}' + a\mathbf{v}' + a'\mathbf{v}]$
- ▶ the cross product is not commutative, so the multiplication of quaternions is not
- ▶ the multiplication of quaternions is
 - ▶ associative
 - ▶ distributive with respect to addition
- ▶ multiplication by a scalar is commutative



Norm of quaternions

- ▶ *conjugation* of $[a, \mathbf{v}]$ is $[a, -\mathbf{v}]$

Norm of quaternions

- ▶ *conjugation* of $[a, \mathbf{v}]$ is $[a, -\mathbf{v}]$
- ▶ conjugation has some natural properties
 - ▶ $\overline{\overline{q}} = q$
 - ▶ $\overline{pq} = \overline{q}\overline{p}$
 - ▶ $\overline{p+q} = \overline{p} + \overline{q}$
 - ▶ $\overline{q}q = q\overline{q}$

Norm of quaternions

- ▶ *conjugation* of $[a, \mathbf{v}]$ is $[a, -\mathbf{v}]$
- ▶ conjugation has some natural properties
 - ▶ $\overline{\overline{q}} = q$
 - ▶ $\overline{pq} = \overline{q}\overline{p}$
 - ▶ $\overline{p+q} = \overline{p} + \overline{q}$
 - ▶ $\overline{q}q = q\overline{q}$
- ▶ *norm of quaternion* $\|q\| = \sqrt{q\overline{q}}$

Norm of quaternions

- ▶ *conjugation* of $[a, \mathbf{v}]$ is $[a, -\mathbf{v}]$
- ▶ conjugation has some natural properties
 - ▶ $\overline{\overline{q}} = q$
 - ▶ $\overline{pq} = \overline{q}\overline{p}$
 - ▶ $\overline{p+q} = \overline{p} + \overline{q}$
 - ▶ $\overline{q}q = q\overline{q}$
- ▶ *norm of quaternion* $\|q\| = \sqrt{q\overline{q}}$
 - ▶ $\|qq'\| = \|q\|\|q'\|$

Norm of quaternions

- ▶ *conjugation* of $[a, \mathbf{v}]$ is $[a, -\mathbf{v}]$
- ▶ conjugation has some natural properties
 - ▶ $\overline{\overline{q}} = q$
 - ▶ $\overline{pq} = \overline{q}\overline{p}$
 - ▶ $\overline{p+q} = \overline{p} + \overline{q}$
 - ▶ $\overline{q}q = q\overline{q}$
- ▶ *norm of quaternion* $\|q\| = \sqrt{q\overline{q}}$
 - ▶ $\|qq'\| = \|q\|\|q'\|$
 - ▶ $q^{-1} = \frac{\overline{q}}{\|q\|^2}$

The map $v \rightarrow q_1 v q_2^{-1}$

- ▶ we already know $\|q_1 v q_2^{-1}\| = \|q_1\| \|v\| \|q_2^{-1}\|$

The map $v \rightarrow q_1 v q_2^{-1}$

- ▶ we already know $\|q_1 v q_2^{-1}\| = \|q_1\| \|v\| \|q_2^{-1}\|$
- ▶ so the map $v \rightarrow q_1 v q_2^{-1}$ is a similarity of 4D space, which multiply every length by $\sqrt{\|q_1\| \|q_2\|}$

The map $v \rightarrow q_1 v q_2^{-1}$

- ▶ we already know $\|q_1 v q_2^{-1}\| = \|q_1\| \|v\| \|q_2^{-1}\|$
- ▶ so the map $v \rightarrow q_1 v q_2^{-1}$ is a similarity of 4D space, which multiply every length by $\sqrt{\|q_1\| \|q_2\|}$
- ▶ in the case $\|q_1\| \|q_2\| = 1$ it is an isometry

The map $v \rightarrow q_1 v q_2^{-1}$

- ▶ we already know $\|q_1 v q_2^{-1}\| = \|q_1\| \|v\| \|q_2^{-1}\|$
- ▶ so the map $v \rightarrow q_1 v q_2^{-1}$ is a similarity of 4D space, which multiply every length by $\sqrt{\|q_1\| \|q_2\|}$
- ▶ in the case $\|q_1\| \|q_2\| = 1$ it is an isometry
- ▶ if $v = 1$ is fixed, also the space (perpendicular to v) of elements $bi + cj + dk$ is

The map $v \rightarrow q_1 v q_2^{-1}$

- ▶ we already know $\|q_1 v q_2^{-1}\| = \|q_1\| \|v\| \|q_2^{-1}\|$
- ▶ so the map $v \rightarrow q_1 v q_2^{-1}$ is a similarity of 4D space, which multiply every length by $\sqrt{\|q_1\| \|q_2\|}$
- ▶ in the case $\|q_1\| \|q_2\| = 1$ it is an isometry
- ▶ if $v = 1$ is fixed, also the space (perpendicular to v) of elements $bi + cj + dk$ is
- ▶ equivalently $v = 1$ is fixed iff $q_1 q_2 = 1$, i. e. $q_1 = q_2^{-1}$

The properties of the map $v \rightarrow qvq^{-1}$

- ▶ the map $v \rightarrow qvq^{-1}$ is isometry of 3D space

The properties of the map $v \rightarrow qvq^{-1}$

- ▶ the map $v \rightarrow qvq^{-1}$ is isometry of 3D space
- ▶ for q and rq ($r \in \mathbb{R}$) is the map same

The properties of the map $v \rightarrow qvq^{-1}$

- ▶ the map $v \rightarrow qvq^{-1}$ is isometry of 3D space
- ▶ for q and rq ($r \in \mathbb{R}$) is the map same
- ▶ we can restrict to *unit quaternions*, which satisfies $\|q\| = 1$

The properties of the map $v \rightarrow qvq^{-1}$


- ▶ the map $v \rightarrow qvq^{-1}$ is isometry of 3D space
- ▶ for q and rq ($r \in \mathbb{R}$) is the map same
- ▶ we can restrict to *unit quaternions*, which satisfies $\|q\| = 1$
- ▶ for q unit quaternion the map preserves the norm of the complex part

The properties of the map $v \rightarrow qvq^{-1}$

- ▶ the map $v \rightarrow qvq^{-1}$ is isometry of 3D space
- ▶ for q and rq ($r \in \mathbb{R}$) is the map same
- ▶ we can restrict to *unit quaternions*, which satisfies $\|q\| = 1$
- ▶ for q unit quaternion the map preserves the norm of the complex part
- ▶ the map commute with multiplication of the complex part by a scalar

Rotation in 3D


We would like to find a formula for rotation of vector $\mathbf{r} \in \mathbb{R}^3$ about vector \mathbf{n} throughout the angle θ .



Rotation in 3D

We would like to find a formula for rotation of vector $\mathbf{r} \in \mathbb{R}^3$ about vector \mathbf{n} throughout the angle θ .


► $\mathbf{r} = \mathbf{r}_1 + \mathbf{r}_2$



Rotation in 3D

We would like to find a formula for rotation of vector $\mathbf{r} \in \mathbb{R}^3$ about vector \mathbf{n} throughout the angle θ .


- ▶ $\mathbf{r} = \mathbf{r}_1 + \mathbf{r}_2$
 - ▶ \mathbf{r}_1 projection to \mathbf{n} , $\mathbf{r}_1 = (\mathbf{r} \cdot \mathbf{n})\mathbf{n}$
 - ▶ \mathbf{r}_2 perpendicular to \mathbf{n} , $\mathbf{r}_2 = \mathbf{r} - \mathbf{r}_1$



Rotation in 3D

We would like to find a formula for rotation of vector $\mathbf{r} \in \mathbb{R}^3$ about vector \mathbf{n} throughout the angle θ .


- ▶ $\mathbf{r} = \mathbf{r}_1 + \mathbf{r}_2$
 - ▶ \mathbf{r}_1 projection to \mathbf{n} , $\mathbf{r}_1 = (\mathbf{r} \cdot \mathbf{n})\mathbf{n}$
 - ▶ \mathbf{r}_2 perpendicular to \mathbf{n} , $\mathbf{r}_2 = \mathbf{r} - \mathbf{r}_1$
- ▶ \mathbf{v} perpendicular to \mathbf{n} , $\mathbf{v} = \mathbf{n} \times \mathbf{r}_2 = \mathbf{n} \times \mathbf{r}$



Rotation in 3D

We would like to find a formula for rotation of vector $\mathbf{r} \in \mathbb{R}^3$ about vector \mathbf{n} throughout the angle θ .


- ▶ $\mathbf{r} = \mathbf{r}_1 + \mathbf{r}_2$
 - ▶ \mathbf{r}_1 projection to \mathbf{n} , $\mathbf{r}_1 = (\mathbf{r} \cdot \mathbf{n})\mathbf{n}$
 - ▶ \mathbf{r}_2 perpendicular to \mathbf{n} , $\mathbf{r}_2 = \mathbf{r} - \mathbf{r}_1$
- ▶ \mathbf{v} perpendicular to \mathbf{n} , $\mathbf{v} = \mathbf{n} \times \mathbf{r}_2 = \mathbf{n} \times \mathbf{r}$
- ▶ $(\mathbf{r}')_2 = \mathbf{r}_2 \cos \theta + \mathbf{v} \sin \theta$
- ▶ $(\mathbf{r}')_1 = \mathbf{r}_1$



Rotation in 3D

We would like to find a formula for rotation of vector $\mathbf{r} \in \mathbb{R}^3$ about vector \mathbf{n} throughout the angle θ .

- ▶ $\mathbf{r} = \mathbf{r}_1 + \mathbf{r}_2$
 - ▶ \mathbf{r}_1 projection to \mathbf{n} , $\mathbf{r}_1 = (\mathbf{r} \cdot \mathbf{n})\mathbf{n}$
 - ▶ \mathbf{r}_2 perpendicular to \mathbf{n} , $\mathbf{r}_2 = \mathbf{r} - \mathbf{r}_1$
- ▶ \mathbf{v} perpendicular to \mathbf{n} , \mathbf{r}_2 , $\mathbf{v} = \mathbf{n} \times \mathbf{r}_2 = \mathbf{n} \times \mathbf{r}$
- ▶ $(\mathbf{r}')_2 = \mathbf{r}_2 \cos \theta + \mathbf{v} \sin \theta$
- ▶ $(\mathbf{r}')_1 = \mathbf{r}_1$
- ▶ $\mathbf{r}' = (1 - \cos \theta)(\mathbf{r} \cdot \mathbf{n})\mathbf{n} + \mathbf{r} \cos \theta + (\mathbf{n} \times \mathbf{r}) \sin \theta$



Rotation in 3D

Now, we find a formula for the map $v \rightarrow qvq^{-1}$.

Rotation in 3D

Now, we find a formula for the map $v \rightarrow qvq^{-1}$.

- ▶ every unit quaternion is in the form $q = [\cos \theta, \mathbf{n} \sin \theta]$, where $|\mathbf{n}| = 1$

Rotation in 3D

Now, we find a formula for the map $v \rightarrow qvq^{-1}$.

- ▶ every unit quaternion is in the form $q = [\cos \theta, \mathbf{n} \sin \theta]$, where $|\mathbf{n}| = 1$
- ▶ we will use $\mathbf{v}_1 \times (\mathbf{v}_2 \times \mathbf{v}_3) = (\mathbf{v}_1 \cdot \mathbf{v}_3)\mathbf{v}_2 - (\mathbf{v}_1 \cdot \mathbf{v}_2)\mathbf{v}_3$

Rotation in 3D

Now, we find a formula for the map $v \rightarrow qvq^{-1}$.

- ▶ every unit quaternion is in the form $q = [\cos \theta, \mathbf{n} \sin \theta]$, where $|\mathbf{n}| = 1$
- ▶ we will use $\mathbf{v}_1 \times (\mathbf{v}_2 \times \mathbf{v}_3) = (\mathbf{v}_1 \cdot \mathbf{v}_3)\mathbf{v}_2 - (\mathbf{v}_1 \cdot \mathbf{v}_2)\mathbf{v}_3$
- ▶ to expand
$$[s, \mathbf{v}][0, \mathbf{r}][s, \mathbf{v}]^{-1} = [0, (s^2 - \mathbf{v} \cdot \mathbf{v} + 2(\mathbf{v} \cdot \mathbf{r})\mathbf{v} + 2s(\mathbf{v} \times \mathbf{r}))]$$

Rotation in 3D

Now, we find a formula for the map $v \rightarrow qvq^{-1}$.

- ▶ every unit quaternion is in the form $q = [\cos \theta, \mathbf{n} \sin \theta]$, where $|\mathbf{n}| = 1$
- ▶ we will use $\mathbf{v}_1 \times (\mathbf{v}_2 \times \mathbf{v}_3) = (\mathbf{v}_1 \cdot \mathbf{v}_3)\mathbf{v}_2 - (\mathbf{v}_1 \cdot \mathbf{v}_2)\mathbf{v}_3$
- ▶ to expand
$$[s, \mathbf{v}][0, \mathbf{r}][s, \mathbf{v}]^{-1} = [0, (s^2 - \mathbf{v} \cdot \mathbf{v} + 2(\mathbf{v} \cdot \mathbf{r})\mathbf{v} + 2s(\mathbf{v} \times \mathbf{r}))]$$
- ▶ substituting $s = \cos \theta$ and $\mathbf{v} = \mathbf{n} \sin \theta$

Rotation in 3D

Now, we find a formula for the map $v \rightarrow qvq^{-1}$.

- ▶ every unit quaternion is in the form $q = [\cos \theta, \mathbf{n} \sin \theta]$, where $|\mathbf{n}| = 1$
- ▶ we will use $\mathbf{v}_1 \times (\mathbf{v}_2 \times \mathbf{v}_3) = (\mathbf{v}_1 \cdot \mathbf{v}_3)\mathbf{v}_2 - (\mathbf{v}_1 \cdot \mathbf{v}_2)\mathbf{v}_3$
- ▶ to expand
$$[s, \mathbf{v}][0, \mathbf{r}][s, \mathbf{v}]^{-1} = [0, (s^2 - \mathbf{v} \cdot \mathbf{v} + 2(\mathbf{v} \cdot \mathbf{r})\mathbf{v} + 2s(\mathbf{v} \times \mathbf{r}))]$$
- ▶ substituting $s = \cos \theta$ and $\mathbf{v} = \mathbf{n} \sin \theta$
- ▶ we have $[s, \mathbf{v}][0, \mathbf{r}][s, \mathbf{v}]^{-1} = [0, (1 - \cos 2\theta)(\mathbf{r} \cdot \mathbf{n})\mathbf{n} + \mathbf{r} \cos 2\theta + (\mathbf{n} \times \mathbf{r}) \sin 2\theta]$

Quaternions and 3D geometry

Theorem (Euler)

The product of two rotation is another rotation.

Quaternions and 3D geometry

Theorem (Euler)

The product of two rotation is another rotation.

Proof.

$$q_2(q_1 p q_1^{-1})q_2^{-1} = (q_2 q_1) p (q_2 q_1)^{-1}$$



Quaternions and 3D geometry

Theorem (Euler)

The product of two rotation is another rotation.

Proof.

$$q_2(q_1 p q_1^{-1})q_2^{-1} = (q_2 q_1)p(q_2 q_1)^{-1}$$



Theorem

The map that take q to map $v \rightarrow qvq^{-1}$ is a 2-to-1 homomorphism from the group of unit quaternions to SO_3 .

Quaternions and 3D geometry

Theorem (Euler)

The product of two rotation is another rotation.

Proof.

$$q_2(q_1 p q_1^{-1})q_2^{-1} = (q_2 q_1)p(q_2 q_1)^{-1}$$



Theorem

The map that take q to map $v \rightarrow qvq^{-1}$ is a 2-to-1 homomorphism from the group of unit quaternions to SO_3 .

Proof.

It is 2-to-1 because q and $-q$ yields the same rotation.



Quaternions and 3D geometry

Theorem (Euler)

The product of two rotation is another rotation.

Proof.

$$q_2(q_1 p q_1^{-1})q_2^{-1} = (q_2 q_1)p(q_2 q_1)^{-1}$$



Theorem

The map that take q to map $v \rightarrow qvq^{-1}$ is a 2-to-1 homomorphism from the group of unit quaternions to SO_3 .

Proof.

It is 2-to-1 because q and $-q$ yields the same rotation.



- unit quaternions is a double cover of SO_3

Higher dimensions

- ▶ John Thomas Graves: *"If with your alchemy you can make three pounds of gold, why should you stop there?"*

Higher dimensions

- ▶ John Thomas Graves: "*If with your alchemy you can make three pounds of gold, why should you stop there?*"
- ▶ 26 December 1843 octaves

Higher dimensions

- ▶ John Thomas Graves: "*If with your alchemy you can make three pounds of gold, why should you stop there?*"
- ▶ 26 December 1843 octaves
- ▶ January 1844 general theory of 2^m -ions, not successful

Higher dimensions

- ▶ John Thomas Graves: *"If with your alchemy you can make three pounds of gold, why should you stop there?"*
- ▶ 26 December 1843 octaves
- ▶ January 1844 general theory of 2^m -ions, not successful
- ▶ In the correspondence of Hamilton and Graves first time the notion of associativity

Higher dimensions

- ▶ John Thomas Graves: *"If with your alchemy you can make three pounds of gold, why should you stop there?"*
- ▶ 26 December 1843 octaves
- ▶ January 1844 general theory of 2^m -ions, not successful
- ▶ In the correspondence of Hamilton and Graves first time the notion of associativity
- ▶ Arthur Cayley : relation between the quaternions and hyper elliptic functions

Higher dimensions

- ▶ John Thomas Graves: *"If with your alchemy you can make three pounds of gold, why should you stop there?"*
- ▶ 26 December 1843 octaves
- ▶ January 1844 general theory of 2^m -ions, not successful
- ▶ In the correspondence of Hamilton and Graves first time the notion of associativity
- ▶ Arthur Cayley : relation between the quaternions and hyper elliptic functions
- ▶ March 1845 paper, one part about octonions - Cayley numbers

Survey

Were quaternions

- ▶ discovered?
- ▶ invented?

Thank you for your attention!