

Homework 3

Finite Element Methods 1

Due date: 17th December 2024

Submit a PDF/scan of the answers to the following questions before the deadline via the *Study Group Roster (Záznamník učitele)* in SIS, or hand-in directly at the practical class on the 17th December 2024


- (1 point) Consider a triangulation \mathcal{T}_h of $\Omega \subset \mathbb{R}^2$ consisting of simplices T with diameter h_T , and define by ϱ_T the diameter of the largest inscribed ball in T . Show that the condition

$$\frac{h_T}{\varrho_T} \leq \sigma, \quad \text{for all } T \in \mathcal{T}_h, \quad (1.1)$$

where the constant σ is independent of T , is equivalent to the condition that all angles in all $T \in \mathcal{T}_h$ are bounded from below by a positive constant θ_0 independent of T .


Solution:

Let T be a triangle and (1.1) be satisfied. Let θ_T be the smallest angle of T , A be the vertex of T corresponding to θ_T , and B, C the other two vertices. Define $b = |AC|$, and v as the distance of C from the line AB . Then,



$$\theta_T \leq \frac{\pi}{3} \quad \text{and} \quad \sin \theta_T = \frac{v}{b} \geq \frac{\varrho_T}{h_T} \geq \frac{1}{\sigma} \quad \implies \quad \theta_T \geq \arcsin \frac{1}{\sigma} =: \theta_0.$$

Therefore, (1.1) \implies all angles bounded from below.



Now assume all angles are bounded from below by $\theta_0 > 0$ and define T' as on the left. Then, for any $T \in \mathcal{T}_h$, T' is contained within T . As the centre of the inscribed circle of T' is at the intersection of the bisectors of the angles, its diameter is given by $h_T \tan \frac{\theta_0}{2} \leq \varrho_T$. Therefore,

$$\frac{h_T}{\varrho_T} \leq \frac{1}{\tan \frac{\theta_0}{2}} =: \sigma \quad \forall T \in \mathcal{T}_h.$$

2. Consider a triangulation \mathcal{T}_h of $\Omega \subset \mathbb{R}^n$ consisting of simplices T with diameter h_T , define by ϱ_T the diameter of the largest inscribed ball in T , and assume that (1.1) holds.

- (a) (1 point) Show that $|T|$, for any $T \in \mathcal{T}_h$, satisfies the condition

$$C_1 h_T^n \leq |T| \leq C_2 h_T^n,$$

where C_1 is a positive constant dependent only on σ and n , and C_2 is a positive constant dependent only on n .

Solution:

Let Ω_n be the volume of the unit ball in \mathbb{R}^n ; then, the volume of a ball with diameter d is $\Omega_n (d/2)^n$. As the volume of a simplex is larger than the volume of its inscribed ball

$$|T| \geq \Omega_n \left(\frac{\varrho_T}{2} \right)^n \geq \underbrace{\frac{\Omega_n}{(2\sigma)^n}}_{C_1} h_T^n. \quad (2.2)$$

Furthermore, as T is contained within any ball with radius h_T and centre in T ,

$$|T| \leq \underbrace{\Omega_n}_{C_2} h_T^n \quad (2.3)$$

- (b) (1 point) Show, for $n = 3$, that any face F of \mathcal{T}_h satisfies the condition.

$$\frac{h_F}{\varrho_F} \leq \sigma.$$

Solution:

Consider any simplex $T \in \mathcal{T}_h$ and let F be a face of T . Let $B \subset T$ be ball with centre c contained within T ; therefore, its diameter is less or equal to ϱ_T . Let p be a plane parallel to F passing through c . Then, $p \cap T$ is a similar (smaller) triangle to F , and the circle $p \cap B$ has the same radius as B ; therefore, $\varrho_F \geq \varrho_T$. As $h_F \leq h_T$ then

$$\frac{h_F}{\varrho_F} \leq \frac{h_T}{\varrho_T} \leq \sigma.$$

- (c) (1 point) Show that $h_T \leq \sigma h_{\tilde{T}}$, for $n = 2, 3$, for any pair of elements $T, \tilde{T} \in \mathcal{T}_h$ sharing an edge.

Solution:

Let ℓ be the length of the edge shared by T and \tilde{T} . Then,

$$h_T \leq \varrho_T \sigma \leq \ell \sigma \leq h_{\tilde{T}} \sigma.$$

3. (2 points) Let \mathcal{T}_h be a triangulation consisting of n -simplices T in \mathbb{R}^n satisfying (1.1) and the assumptions (\mathcal{T}_h1) – (\mathcal{T}_h5) . Prove that the number of elements of \mathcal{T}_h sharing a vertex is bounded by a constant depending on σ and n .

Solution:

Let a be a vertex of \mathcal{T}_h and let $\mathcal{M}_a = \{T \in \mathcal{T}_h : a \in T\}$ be the set of simplices sharing the vertex a . Consider any $T \in \mathcal{M}_a$ with vertices a_1, \dots, a_{n+1} , and let \tilde{T} be a simplex with vertices $\tilde{a}_1, \dots, \tilde{a}_{n+1}$ satisfying $(a_i - a) = h_T(\tilde{a}_i - a)$, $i = 1, \dots, n+1$. Then, $\tilde{T} = \phi(T)$, where $\phi(x) = (x-a)/h_T + a$, and $h_{\tilde{T}} = 1$. For a ball $B \subset T$, then $\tilde{B} := \phi(B)$ is a ball in \tilde{T} and $\text{diam}(B) = h_T \text{diam } \tilde{B}$; therefore,

$$\sigma \geq \frac{h_T}{\varrho_T} = \frac{h_T}{h_T \varrho_{\tilde{T}}} = \frac{h_{\tilde{T}}}{\varrho_{\tilde{T}}}.$$

Denoting by Ω_n the volume of the unit ball in \mathbb{R}^n , we have that

$$\Omega_n \geq \sum_{T \in \mathcal{M}_a} |\tilde{T}| \geq \sum_{T \in \mathcal{M}_a} \left(\frac{\varrho_{\tilde{T}}}{2}\right)^n \Omega_n \geq \sum_{T \in \mathcal{M}_a} \left(\frac{1}{2\sigma}\right)^n \Omega_n = \left(\frac{1}{2\sigma}\right)^n \Omega_n \text{card } \mathcal{M}_a;$$

therefore, $\text{card } \mathcal{M}_a \leq (2\sigma)^n$.