

Prop. XIII. 16 $T \in U(H)$ normal, E_T its spectral measure
 $\Rightarrow \tilde{g}(T) = \int g dE_T$ for each bdd L^2 -meas. function g

Proof: $x, y \in H \Rightarrow \langle \tilde{g}(T)x, y \rangle = \int \underset{\uparrow}{g} d(E_T)_{xy} = \underset{\substack{\text{construction} \\ \text{of meas.-calculus}}}{\langle \int g dE_T \rangle_{xy}}$ $\underset{\uparrow}{= \langle \int g dE_T \rangle_{xy}}$ $\underset{\substack{\text{defn. of } \int g dE_T}}{= \langle \int g dE_T \rangle_{xy}}$

Lemma VI.15 T self-adjoint (unbdd), C_T its Cayley transform, E ... the spectral measure of C_T

$$\text{Then } T = \int c \frac{1+z}{1-z} dE(z)$$

Proof ① C_T is unitary $\Rightarrow \sigma(C_T) \subset \pi = \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}$

② $I - C_T$ is one-to-one, i.e. 1 is not an eigenvalue of $C_T \Rightarrow E(\{1\}) = 0$ [by Ax. 10 & Prop. 13]

③ $f(z) = c \cdot \frac{1+z}{1-z}$ is \mathbb{R} -measurable (defined E -a.e.)

$$\begin{aligned} f \text{ is real-valued} \quad \dots \quad c \frac{1+z}{1-z} &= c \frac{(1+z)(1-\bar{z})}{(1+z)(1-\bar{z})} = \\ (\text{essentially}) \quad &= c \frac{1+z-\bar{z}-|z|^2}{(1-z)\bar{z}} = -\frac{z \operatorname{Im} z}{|z-1|^2} \\ |z|=1, z \in \pi \end{aligned}$$

$\Rightarrow S := \int f dE$ is self-adjoint (Thm 12 (c))

④ Moreover: $f(z) \circ (I-z) = c \cdot (1+z)$

$$\text{so, by Thm 12 (s): } S(I - C_T) = c(I + C_T)$$

$$\begin{aligned} \Rightarrow S(I - C_T)(I - C_T)^{-1} &= c(I + C_T)(I - C_T)^{-1} \\ I \not\models D(C_T) &\qquad \qquad \qquad T \text{ (Thm V.3 (c))} \\ \{R(I - C_T) = D(T)\} &\\ \text{Thm V.3 (s)} \end{aligned}$$

$$\Rightarrow S \models D(T) = \overline{T} \Rightarrow T \subset S$$

S, T self-adjoint $\Rightarrow T = S$

Lemma III.16

F abstract spectral measure or at

$\varphi: \mathcal{A} \rightarrow \mathbb{C}$ \mathcal{B} -measurable

$$E(A) = F(\varphi^{-1}(A)), \quad A \in \mathcal{B}' = \{A \subset \mathbb{C}, \varphi^{-1}(A) \subset \mathcal{A}\}$$

(1) E is an abstract spectral measure

• properties (i) - (vii) are obvious

• (viii): $E_{x,y}$ is a complex Borel measure for each $x, y \in H$
By remark after 26 it is enough to prove it for
 $E_{x,x} \quad x \in H$

Note that $E_{x,x} = \varphi(F_{x,x})$. To simplify notation

$$\text{set } \mu := F_{x,x}, \quad \nu := E_{x,x} = \varphi(\mu)$$

Let $A \in \mathcal{B}'$ set $\beta := \sup \{\nu(B), B \subset A \text{ Borel}\}$
 $\gamma := \inf \{\nu(C), C \supset A \text{ Borel}\}$

Let $B_n \subset A \subset C_n$ be Borel s.t. $\nu(B_n) > \beta - \frac{1}{n}$
 $\nu(C_n) < \gamma + \frac{1}{n}$

$$B := \bigcup_n B_n, \quad C := \bigcap_n C_n \Rightarrow B, C \text{ Borel}, \quad B \subset C \subset \mathcal{C}$$

 $\nu(B) = \beta, \quad \nu(C) = \gamma$

We will be done if we prove $\nu(C \setminus B) = 0$ (i.e. $\beta = \gamma$)

Suppose $\nu(C \setminus B) > 0$. Then either $\nu(C \setminus A) > 0$ or $\nu(A \setminus B) > 0$

Suppose $\nu(C \setminus A) > 0$ (+ the other case is analogous)

Then $\mu(\varphi^{-1}(C \setminus A)) = \nu(C \setminus A) > 0, \quad \varphi^{-1}(C \setminus A) \in \mathcal{A}$

Since μ is a Borel measure, there is $D \subset \varphi^{-1}(C \setminus A)$

Borel s.t. $\mu(D) > 0$ [in fact $\mu(D) = \mu(\varphi^{-1}(C \setminus A))$]

Since Borel measures on \mathbb{C} are regular, there is $D_n \subset D$ compact
s.t. $\mu(D_n) > 0$

By Luzin's theorem there is $k \in D$, compact s.t. $\mu(k) > 0$
 & $\varphi|_K$ is continuous

Then $\varphi(k) \subset C \setminus A$, $\varphi(k)$ is compact, hence Borel,
 and $\nu(\varphi(k)) = \mu(\varphi^{-1}(\varphi(k))) \geq \mu(k) > 0$

So, $C \setminus \varphi(k) \supset A$ is a Borel set with $\nu(C \setminus \varphi(k)) < \gamma$,
 a contradiction completing the proof.

(2) $f: C \rightarrow \mathbb{C}$ \mathcal{F}^1 -measurable $\Rightarrow \int f dE = \int (f \circ \varphi) dF$

• $\int |f|^2 dE_{+,+} = \int |f|^2 d\mu_{F_{+,+}} = \int |f \circ \varphi|^2 dF_{+,+}$

\Rightarrow the two domains coincide

• $x, y \in D (\int f dE) \Rightarrow$

$$\langle (\int f dE)_{+,y} \rangle = \int f dE_{+,y} = \int f d\mu_{F_{+,y}} = \int f \circ \varphi dF_{+,y}$$

$$= \langle (\int (f \circ \varphi) dF)_{+,y} \rangle$$

Theorem XIII.17 T self-adjoint $\Rightarrow \exists!$ abstract spectral measure
 E s.t. $T = \int c dE$

Moreover, this E is the image of the spectral measure of C_T
under $z \mapsto c \frac{1+z}{1-z}$

Proof ① Let F be the spectral measure of C_T

$$\varphi(z) = c \frac{1+z}{1-z}, z \in \mathbb{C} \setminus \{\pm i\}$$

Since $F(\{\pm i\}) = 0$, φ is measurable (see the proof of Lemma 15)

Let $E = \varphi(F)$. By L 16, E is an abstract spectral measure and

$$\int c dE = \int c d\varphi dF = \int \varphi dF = T$$

② Uniqueness: Let E be an abstract spectral measure such that $T = \int c dE$

$$T(C_T) \subset \mathbb{R} \Rightarrow \text{ess-supp}(cd) \subset \mathbb{R}$$

$$\text{Set } g(z) = \frac{z-c}{z+c} \quad \text{for } z \in \mathbb{R} \text{ we have } |g(z)| = 1$$

$$\frac{1}{g(z)} = \overline{g(z)}$$

$\Rightarrow U := \int g dE$ is a unitary operator

$$\text{By Prop. 14} \quad E_U = g(E)$$

$$\text{Further, } \varphi \circ g = cd \quad E-a.s. \Rightarrow$$

$$\Rightarrow T = \int cd dE = \int \varphi \circ g dE = \int \varphi dE_U$$

$$\Rightarrow T(I - U) = (\int \varphi dE_U)(I - U) = (\int \varphi dE_U)(\int (1-z) dE_U(z))$$

Thm 12(3)

$$= \int c(1+z) dE_U(z) = c(I + U) \Rightarrow U = CT$$

Since $E = \varphi(E_u)$, we deduce the uniqueness.

Corollary XIII.18 T self-adjoint $\Rightarrow (T \text{ bdd} \Leftrightarrow \sigma(T) \text{ bdd})$

Proof \Rightarrow clear, spectrum of a bdd operator is compact

$\Leftarrow T = cd \text{ dE}$, $E(\sigma \cap \sigma(T)) = 0$ and
 $\sigma(T) \text{ bdd} \Rightarrow cd$ is essentially bdd